

LIE BIALGEBRAS AND THE CYCLIC HOMOLOGY OF A_∞ STRUCTURES IN TOPOLOGY

XIAOJUN CHEN

ABSTRACT. A_∞ categories are a mathematical structure that appears in topological field theory, string topology, and symplectic topology. This paper studies the cyclic homology of a Calabi-Yau A_∞ category, and shows that it is naturally an equivariant topological conformal field theory, and in particular, contains an involutive Lie bialgebra. Applications of the theory to string topology and the Fukaya category are given; in particular, it is shown that there is a Lie bialgebra homomorphism from the cyclic cohomology of the Fukaya category of a symplectic manifold with contact type boundary to the linearized contact homology of the boundary.

CONTENTS

1. Introduction	1
2. Cyclic homology of A_∞ categories	3
2.1. Examples from topology	6
3. Calabi-Yau A_∞ categories	9
3.1. Topological conformal field theory	9
4. The Lie bialgebra structure	12
4.1. Construction of the Lie bialgebra	12
4.2. Noncommutative symplectic geometry	17
4.3. Example from string topology	19
5. Application in symplectic topology	20
5.1. Symplectic field theory	20
5.2. From Fukaya category to symplectic field theory	22
References	23

1. INTRODUCTION

In this article we study some algebraic structures on the cyclic homology of an A_∞ category that appears in topology. A_∞ structures have been widely studied in the past decades by mathematicians from various fields, such as symplectic topology, homological algebra, and string field theory. On the other hand, the cyclic homology of algebras and categories has also played an important role in the study of, for example, the non-commutative geometry and K -theory. The motivation for us to study the cyclic homology of A_∞ structures is the recent work of Costello ([10]), which asserts that:

- A Calabi-Yau A_∞ category $\mathcal{A}$ is homotopy equivalent to an open topological conformal field theory (TCFT);
- Associated to a Calabi-Yau A_∞ category $\mathcal{A}$ is a natural open-closed TCFT;

- The closed states of the associated open-closed TCFT is the Hochschild cohomology of $\mathcal{A}$.

Some similar work has also previously been obtained by Kontsevich, part of which was summarized in [27, §11]. A Calabi-Yau A_∞ category is an A_∞ category with a non-degenerate pairing on the morphism spaces. Among the examples are the Fukaya category of Lagrangian submanifolds of symplectic manifolds, and the category of coherent sheaves on Calabi-Yau manifolds. Roughly speaking, a *closed* TCFT is a functor from Segal's category to the category of chain complexes, where Segal's category is the category whose objects are the natural numbers (identified with circles) and the morphisms between two sets of circles are (the chain complex of) the moduli space of Riemann surfaces connecting them. The open and open-closed TCFT's are defined similarly, where the objects are the intervals and the unions of intervals and circles respectively.

There is an S^1 action by rotation on the inputs and outputs of a closed TCFT. In [16] Getzler first studied the TCFT's which are invariant under the S^1 action. We shall call these TCFT's *equivariant* (with respect to the S^1 action). One of the key observations is that Costello's functor from the open TCFT to the closed TCFT respects the S^1 rotation. By modulo such a rotation, one obtains on the one side, the cyclic cohomology of a Calabi-Yau A_∞ category; and on the other side, an equivariant TCFT:

Theorem 1 (Theorem 6). *Let $\mathcal{A}$ be a Calabi-Yau A_∞ category. Then the cyclic cohomology of $\mathcal{A}$ has the structure of an equivariant topological conformal field theory.*

In [15, 16] Getzler showed that a closed TCFT has the structure of a Batalin-Vilkovisky algebra while an equivariant TCFT has the structure of a *gravity algebra*. A gravity algebra is a Lie algebra with more constraints, which he called the *generalized Jacobi identities*. We shall give an explicit formula for the Lie algebra of Getzler on the cyclic cohomology of a Calabi-Yau A_∞ category. In fact, we show more, namely, the cyclic cohomology is an *involutive Lie bialgebra* (see Def. 7):

Theorem 2 (Theorem 7). *Let $\mathcal{A}$ be a Calabi-Yau A_∞ category. Then the cyclic cochain complex of $\mathcal{A}$ has the structure of a differential graded involutive Lie bialgebra.*

The construction of the Lie bialgebra is inspired by string topology ([5]), and a similar result has been obtained by Barannikov ([3, §5]) and Kontsevich-Soibelman ([27, §11]) in a slightly different and conceptual manner. In string topology, Chas and Sullivan showed that the equivariant homology of the free loop space of a compact manifold, say M , also endows a Lie bialgebra structure. To compare these two structures, recall that K.-T. Chen ([6]) and J. D. S. Jones ([22]) showed that the cyclic cohomology of the cochain algebra of a simply connected M is isomorphic to the equivariant homology of its free loop space LM . However, a transversality issue in string topology obstructs one to model the Lie bialgebra of string topology on the cyclic cohomology of *singular* or *cellular* cochain algebra of M . With the great efficiency of rational homotopy theory, Hamilton-Lazarev ([21]) and Lambrechts-Stanley ([28]) independently construct a cyclic A_∞ algebra over $\mathbb{Q}$, which models the cochain algebra of M with Poincaré duality (in fact the one of Lambrechts-Stanley is a differential graded algebra). Recall that a cyclic A_∞ algebra is a Calabi-Yau A_∞ category with one object, and one has the following:

Theorem 3 (Theorem 10). *Suppose M is a simply connected, compact manifold and V is the cyclic A_∞ algebra which models M , then the Lie bialgebra on the cyclic cohomology of V models that of Chas and Sullivan in string topology.*

Another application of Theorems 1 and 2 is to symplectic topology. Recall that the Fukaya category of a compact symplectic manifold (with or without boundary; for the former case, some convexity conditions are required, for example, the boundary is of contact type as in Example 2 in §2.1) is a Calabi-Yau A_∞ category ([12, 13]). A direct consequence of the above theorems is:

Corollary 1. *The cyclic cohomology of the Fukaya category of a symplectic manifold is naturally an equivariant topological conformal field theory, and in particular, it has the structure of an involutive Lie bialgebra.*

As an application to this corollary, we give a relationship between the Fukaya category and symplectic field theory introduced by Eliashberg, Givental and Hofer ([11]). Suppose M is a symplectic manifold with contact type boundary W . Mathematicians have obtained many interesting structures for M and W in symplectic field theory; in particular, Cieliebak and Latschev showed in [8] that, the linearized contact homology of W (a homology theory on the space of Reeb orbits in W) endows the structure of an involutive Lie bialgebra. Inspired by their work, we show that:

Theorem 4 (Theorem 13). *Let M be an exact symplectic manifold with contact type boundary W . There is a chain map from the linearized contact chain complex of W to the cyclic cochain complex of the Fukaya category of M , which induces on homology a map of Lie bialgebras.*

The paper is organized as follows: In Section 2, we recall the definitions of an A_∞ category and its cyclic homology; some examples arising from topology are given. In Section 3 we give the definition of a Calabi-Yau category and discuss its role in the study of topological conformal field theories. This proves Theorem 1. In Section 4 we first prove Theorem 2. Some relationship between the Lie bialgebras on the cyclic chain complex of an A_∞ algebra and in the non-commutative symplectic geometry is also discussed. Examples from string topology are given and Theorem 3 is shown. In Section 5 we discuss some symplectic field theory and prove Theorem 4.

Acknowledgements. The current work is inspired by the talks given by Costello and Latschev at the CUNY Einstein Chair Seminar. The author also benefits from interesting conversations with Sullivan and Fukaya. He is very grateful to Yongbin Ruan for many encouragements during the preparation of the paper.

2. CYCLIC HOMOLOGY OF A_∞ CATEGORIES

Definition 1 (A_∞ category). Let k be a field. An A_∞ category $\mathcal{A}$ consists of a set of objects $\mathcal{O}b(\mathcal{A})$, a graded k -vector space $\text{Hom}(A_1, A_2)$ for each pair of objects $A_1, A_2 \in \mathcal{O}b(\mathcal{A})$, and a sequence of operators:

$$m_i : \text{Hom}(A_0, A_1) \otimes \text{Hom}(A_1, A_2) \otimes \cdots \otimes \text{Hom}(A_{i-1}, A_i) \rightarrow \text{Hom}(A_0, A_i),$$

where $|m_i| = i - 2$, for $i = 1, 2, \dots$, satisfying the following A_∞ relations:

$$\sum_{p=0}^{i-1} \sum_{k=1}^{i-p} (-1)^{|a_1| + \cdots + |a_p| + p + \epsilon_{i-k+1} + \epsilon_k} m_{i-k+1}(a_1, \dots, a_p, m_k(a_{p+1}, \dots, a_{p+k}), a_{p+k+1}, \dots, a_i) = 0,$$

where

$$\epsilon_k = (k-1)|a_{p+1}| + (k-2)|a_{p+2}| + \cdots + |a_{p+k-1}| + \frac{k(k-1)}{2}$$

and

$$\begin{aligned} \epsilon_{i-k+1} = & (i-k)|a_1| + \cdots + (i-k-p+1)|a_p| + (i-k-p)(i-2+|a_{p+1}| + \cdots + |a_{p+k}|) \\ & + (i-k-p-1)|a_{p+k+1}| + \cdots + |a_{i-1}| + \frac{(i-k)(i-k-1)}{2}. \end{aligned}$$

An A_∞ category with one object is an A_∞ algebra in the usual sense, and conversely an A_∞ category may be viewed as an A_∞ algebra “with several objects”. If an A_∞ category $\mathcal{A}$ has vanishing m_i , for $i = 3, 4, \dots$, then it is a differential graded (DG) category.

Convention. (1) The signs are always complicated in the definition of A_∞ structures. First, recall the Koszul sign convention: Suppose V is a graded space, and $\alpha, \beta \in V$; then in any case when we switch $\alpha \otimes \beta$ to $\beta \otimes \alpha$, there is a $(-1)^{|\alpha||\beta|}$ popping out, i.e., $\alpha \otimes \beta \mapsto (-1)^{|\alpha||\beta|} \beta \otimes \alpha$.

(2) Suppose $(V, \{m_i\})$ is an A_∞ algebra. Let $\Sigma V = V[1]$ be the desuspension of V , and denote by $T(\Sigma V)$ the cotensor algebra generated by ΣV . Define

$$\overline{m}_i = \Sigma \circ m_i \circ (\Sigma^{-1})^{\otimes i} : (\Sigma V)^{\otimes i} \rightarrow \Sigma V, \quad i = 1, 2, \dots$$

i.e., if we denote the identity $id : V \rightarrow \Sigma V, a \mapsto \overline{a}$, then

$$\overline{m}_i(\overline{a}_1, \dots, \overline{a}_i) = (-1)^{(i-1)|\overline{a}_1| + (i-2)|\overline{a}_2| + \cdots + |\overline{a}_{i-1}|} \Sigma(m_i(a_1, a_2, \dots, a_i)).$$

Then $\overline{m} = \overline{m}_1 + \overline{m}_2 + \cdots$ defines a degree -1 co-differential on the coalgebra $T(\Sigma V)$. $(T(\Sigma V), \overline{m})$ is called the *bar construction* of V .

Definition 2 (Hochschild homology). Let $\mathcal{A}$ be an A_∞ category. The *Hochschild chain complex* $\text{Hoch}_*(\mathcal{A})$ of $\mathcal{A}$ is the chain complex whose underlying vector space is

$$\bigoplus_{n=1}^{\infty} \bigoplus_{A_0, A_1, \dots, A_n \in \text{Ob}(\mathcal{A})} \text{Hom}(A_0, A_1) \otimes \text{Hom}(A_1, A_2) \otimes \cdots \otimes \text{Hom}(A_n, A_0)$$

with differential b defined by

$$\begin{aligned} & b(a_0, a_1, \dots, a_n) \\ = & \sum_{j=0}^n \sum_{i=0}^{n-j} \pm (m_{i+j+1}(a_{n-j+1}, \dots, a_n, a_0, \dots, a_i), a_{i+1}, \dots, a_{n-j}) \end{aligned} \quad (1)$$

$$+ \sum_{k=0}^{n-1} \sum_{i=1}^{n-k} \pm (a_0, a_1, \dots, a_k, m_i(a_{k+1}, \dots, a_{k+i}), a_{k+i+1}, \dots, a_n), \quad (2)$$

where the signs $\pm$ in (1) and (2) are given below. The associated homology is called the *Hochschild homology* of $\mathcal{A}$, and is denoted by $HH_*(\mathcal{A})$.

With the sign convention discussed in the previous paragraph, the sign in (1) and (2) are assigned as follows: for simplicity, we assume $\mathcal{A}$ has one object A (the general case is completely the same), and let $V = \text{Hom}(A, A)$; the Hochschild complex of $\mathcal{A}$ is nothing but the *twisted tensor product* of V and $T(\Sigma V)$, from which the sign follows. More precisely, for $(a_0, \overline{a}_1, \overline{a}_2, \dots, \overline{a}_n) \in V \otimes T(\Sigma V)$,

$$\begin{aligned} & b(a_0, \overline{a}_1, \overline{a}_2, \dots, \overline{a}_n) \\ = & \sum_j \sum_i (-1)^{\epsilon_1} (\Sigma^{-1} \circ \overline{m}_{i+j+1}(\overline{a}_{n-j+1}, \dots, \overline{a}_n, \overline{a}_0, \overline{a}_1, \dots, \overline{a}_i), \overline{a}_{i+1}, \dots, \overline{a}_{n-j}) \end{aligned}$$

$$+ \sum_k \sum_i (-1)^{\epsilon_2} (a_0, \bar{a}_1, \dots, \bar{a}_k, \bar{m}_i(\bar{a}_{k+1}, \dots, \bar{a}_{k+i}), a_{k+i+1}, \dots, \bar{a}_n),$$

where $\epsilon_1 = (|\bar{a}_{n-j+1}| + \dots + |\bar{a}_n|)(|\bar{a}_0| + \dots + |\bar{a}_{n-j}|)$ and $\epsilon_2 = |a_0| + |\bar{a}_1| + \dots + |\bar{a}_k|$ (note ϵ_1 and ϵ_2 come from the Koszul sign convention). Let id be the identity map:

$$\begin{aligned} id: \quad V \otimes V \otimes \dots \otimes V &\xrightarrow{id \otimes \Sigma^{\otimes n}} V \otimes \Sigma V \otimes \dots \otimes \Sigma V \\ a_0 \otimes a_1 \otimes \dots \otimes a_n &\longmapsto (-1)^{n|a_0| + (n-1)|a_1| + \dots + |a_{n-1}|} a_0 \otimes \bar{a}_1 \otimes \dots \otimes \bar{a}_n. \end{aligned}$$

The sign in (1) and (2) are assigned such that the following diagram commutes:

$$\begin{array}{ccc} V \otimes TV & \xrightarrow{id} & V \otimes T(\Sigma V) \\ \downarrow b & & \downarrow b \\ V \otimes TV & \xrightarrow{id} & V \otimes T(\Sigma V). \end{array}$$

Remark 1. *The definition of the Hochschild homology of an A_∞ algebra was first introduced by Getzler and Jones in [17]. The signs in the above definition are consistent with theirs (see [17, Lemma 1.3 and Definition 3.5]). In that paper the authors also introduced the cyclic homology of an A_∞ algebra, and hence the cyclic cohomology is also defined. While in the paper of Penkava and Schwarz [30], another version of cyclic cohomology is given. An argument similar to Loday [29, Theorem 2.1.5] shows that these two definitions coincide if the ground field k contains $\mathbb{Q}$. The following definition is due to Penkava and Schwarz.*

Definition 3 (Cyclic homology). Suppose $\mathcal{A}$ is an A_∞ category. Let

$$\begin{aligned} t_n : \text{Hom}(A_0, A_1) \otimes \dots \otimes \text{Hom}(A_{n-1}, A_n) \otimes \text{Hom}(A_n, A_0) \\ \rightarrow \text{Hom}(A_n, A_0) \otimes \text{Hom}(A_0, A_1) \otimes \dots \otimes \text{Hom}(A_{n-1}, A_n), \end{aligned}$$

$n = 0, 1, 2, \dots$, be the linear map

$$t_n(a_0, a_1, \dots, a_n) := (-1)^{n+|a_n|(|a_0| + \dots + |a_{n-1}|)} (a_n, a_0, \dots, a_{n-1}), \quad (3)$$

where the factor $(-1)^n$ is the sign of the cyclic operator t_n . Extending t_n to $\text{Hoch}_*(\mathcal{A})$ trivially, and let

$$t = t_0 + t_1 + t_2 + \dots,$$

the cokernel $\text{Hoch}_*(\mathcal{A})/(1-t)$ of $1-t$ forms a chain complex with the induced differential from the Hochschild complex (still denoted by b). Such chain complex is denoted by $\text{Cycl}_*(\mathcal{A})$, and is called the *Connes cyclic complex* of $\mathcal{A}$. Its homology is called the *cyclic homology* of $\mathcal{A}$, and is denoted by $CH_*(\mathcal{A})$.

The *cyclic cohomology* of $\mathcal{A}$ is the homology of the dual cochain complex $\text{Cycl}^*(\mathcal{A})$ of $\text{Cycl}_*(\mathcal{A})$. Namely, suppose $f \in \text{Hom}(\text{Hoch}_*(\mathcal{A}), k)$, then $f \in \text{Cycl}^*(\mathcal{A})$ if and only if for all $\alpha \in \text{Hoch}_*(\mathcal{A})$, $f(\alpha) = f(t(\alpha))$.

The well-definedness of the above definition is similar to the case of the cyclic homology of differential graded algebras (see, for example, Loday [29, §2.1.4]), the proof of which is therefore omitted. Like the Hochschild chain complex case, the signs in the cyclic chain complex are also delicate. An elegant treatment of the signs, which is due to Quillen ([31]), greatly simplifies the issue. As before,

we assume the A_∞ category $\mathcal{A}$ has one object A , and denote the morphism space by V . Let id be the identity map:

$$\begin{aligned} id : \quad V \otimes V \otimes \cdots \otimes V &\xrightarrow{\Sigma^{\otimes n+1}} \Sigma V \otimes \Sigma V \otimes \cdots \otimes \Sigma V \\ a_0 \otimes a_1 \otimes \cdots \otimes a_n &\longmapsto (-1)^{n|a_0|+(n-1)|a_1|+\cdots+|a_{n-1}|} \bar{a}_0 \otimes \bar{a}_1 \otimes \cdots \otimes \bar{a}_n, \end{aligned}$$

and let $\bar{t}$ be the isomorphism

$$\begin{aligned} \bar{t} : \quad \Sigma V \otimes \Sigma V \otimes \cdots \otimes \Sigma V &\longrightarrow \Sigma V \otimes \Sigma V \otimes \cdots \otimes \Sigma V \\ \bar{a}_0 \otimes \bar{a}_1 \otimes \cdots \otimes \bar{a}_n &\longmapsto (-1)^{|\bar{a}_n|(|\bar{a}_0|+\cdots+|\bar{a}_{n-1}|)} \bar{a}_n \otimes \bar{a}_0 \otimes \cdots \otimes \bar{a}_{n-1}. \end{aligned}$$

Note that the sign in the image of $\bar{t}$ is only coming from the Koszul convention. We have the following:

Lemma 1. *Let $(V, \{m_i\})$ be an A_∞ algebra, and let $\bar{m}_i$ and $id, t, \bar{t}$ be as above. The following diagram commutes:*

$$\begin{array}{ccc} TV & \xrightarrow{t, m_i} & TV \\ \downarrow id & & \downarrow id \\ T(\Sigma V) & \xrightarrow{\bar{t}, \bar{m}_i} & T(\Sigma V). \end{array}$$

The proof is a direct check. By the lemma one may identify the cyclic chain complex $\text{Cycl}_*(V)$, with degree shifted down by 1, with $T^+(\Sigma V)/(1 - \bar{t})$, where $T^+(\Sigma V)$ is the augmented ideal of $T(\Sigma V)$, i.e. $T^+(\Sigma V) = T(\Sigma V)/k$. In other words, the cyclic complex of V , up to a degree shift, is isomorphic to space of cyclically coinvariant elements in the augmented bar construction.

For a general A_∞ category $\mathcal{A}$, we apply the same identification. The advantage is that we do not need to care about the signs any more; all the signs, either coming from the A_∞ structure or from the cyclic operator, just follows the Koszul sign convention.

2.1. Examples from topology. The structure of A_∞ algebras and categories naturally arise in topology. Besides the classical example of A_∞ spaces discovered by Stasheff, there are two examples which have been studied in recent years. One is the A_∞ algebra model of a simply-connected smooth manifold, the other is the A_∞ category of Lagrangian submanifolds (the Fukaya category). In these two cases the associated A_∞ categories bear some additional structure, which make them into the so-called *Calabi-Yau A_∞ categories*. We shall discuss the first example in §4.3 and the second in §5.

In the following we present two simplified examples. They are the derived category of complex of coherent sheaves on a Fano toric manifold and the Fukaya-Seidel category of a symplectic Lefschetz fibration. In both cases, some of the A_∞ operators vanish, which therefore greatly simplifies the computation. Despite of the simplicity, they are proved to be an important role in the understanding of the homological mirror symmetry conjecture ([1, 2]).

Example 1 (Derived category of coherent sheaves). Recall that a *triangulated category* is an additive category equipped with a shifting functor and a class of distinguished triangles, satisfying some conditions (see, for example, [14, Chapter IV]). Suppose $\mathcal{K}$ is a triangulated category. An ordered set of objects $E = (E_0, E_1, \cdots, E_n)$ is called a *strong exceptional collection* if

$$\text{Hom}(E_j, E_i[k]) = \begin{cases} k \cdot id, & \text{if } i = j \text{ and } k = 0, \\ 0, & \text{if } j > i \text{ or } k \neq 0. \end{cases} \quad (4)$$

A strong exceptional collection E is called *full* if it generates the category $\mathcal{K}$, that is, the smallest triangulated category that contains E is $\mathcal{K}$ itself.

Lemma 2. *Suppose $\mathcal{K}$ is a triangulated category, and $(E_1, E_2, \dots, E_n)$ is a strong exceptional collection. Denote by B the endomorphism algebra of*

$$E_1 \oplus E_2 \oplus \dots \oplus E_n;$$

then the cyclic homology $CH_(B)$ is isomorphic to $CH_*(k)^{\oplus n+1}$.*

Proof. Due to Equation (4), its cyclic chain complex is exactly

$$\bigoplus_{i=0}^n \bigoplus_{j=1}^{\infty} \underbrace{(\overline{id}_{E_i}, \overline{id}_{E_i}, \dots, \overline{id}_{E_i})}_j / (1 - \bar{t}),$$

and the lemma follows. $\square$

One may view the set $\{E_0, E_1, \dots, E_n\}$ as objects of a category, where $\text{Hom}(E_i, E_j) = \text{Hom}_{\mathcal{K}}(E_i, E_j)$, then the cyclic homology of B is also the cyclic homology of that category.

The above lemma can be used to compute the cyclic homology of some projective varieties. For example, for $\mathbb{CP}^n$, $(\mathcal{O}_{\mathbb{CP}^n}, \mathcal{O}_{\mathbb{CP}^n}(-1), \dots, \mathcal{O}_{\mathbb{CP}^n}(-n))$ forms a full strong exceptional collection for the bounded derived category of coherent sheaves $\mathcal{D}^b(\text{coh}(\mathbb{CP}^n))$. Denote by $\mathcal{E} := \mathcal{O}_{\mathbb{CP}^n} \oplus \mathcal{O}_{\mathbb{CP}^n}(-1) \oplus \dots \oplus \mathcal{O}_{\mathbb{CP}^n}(-n)$; Bondal and Kapranov showed in [4] that there is an equivalence

$$\mathbf{R}\text{Hom}(\mathcal{E}, -) : \mathcal{D}^b(\text{coh}(\mathbb{CP}^n)) \rightarrow \mathcal{D}^b(\text{Mod}(B)),$$

where $\mathcal{D}^b(\text{Mod}(B))$ is the derived category of complexes of right B -modules whose cohomology is finitely generated. Hence, by the above argument, the cyclic homology of $\mathbb{CP}^n$ is isomorphic to $CH_*(k)^{\oplus n+1}$. For some more examples, see [23, 25, 1, 2].

Example 2 (The Fukaya-Seidel category). We briefly recall for non-experts the Fukaya-Seidel category of symplectic Lefschetz fibrations. The construction is due to P. Seidel ([32, 33]).

Recall that an *exact symplectic manifold with contact type boundary* is a quadruple (M, ω, η, J) , where M is a compact $2n$ dimensional manifold with boundary, ω is a symplectic 2-form on M , η is a 1-form such that $\omega = d\eta$ and J is a ω -compatible almost complex structure. These data also satisfy the following two convexity conditions:

- The negative Liouville vector field defined by $\omega(\cdot, X_\eta) = \eta$ points strictly inwards along the boundary of M ;
- The boundary of M is weakly J -convex, which means that any J -holomorphic curves cannot touch the boundary unless they are completely contained in it.

An n dimensional submanifold $L \subset M$ is called *Lagrangian* if $\omega|_L = 0$. We always assume L is closed and is disjoint from the boundary of M . L is called *exact* if $\eta|_L$ is an exact 1-form.

Take k to be a field of characteristic 2. According to symplectic geometry (see Fukaya [12] or Seidel [34]), the exact Lagrangian submanifolds of M form the objects of an A_∞ category, where the morphisms between L_i and L_j are the Floer cochain complex $CF^*(L_i, L_j; k)$, which is spanned by the transversal intersection points of L_i and L_j . The A_∞ operators $\{m_i\}$ are given by counting the

J -holomorphic disks. Namely, for $(p_1, p_2, \dots, p_i) \in \text{Hom}(A_0, A_1) \otimes \dots \otimes \text{Hom}(A_{i-1}, A_i)$,

$$m_i(p_1, p_2, \dots, p_i) = \sum_{q \in \text{Hom}(A_1, A_i)} \# \mathcal{M}(p_1, p_2, \dots, p_i, q) \cdot q,$$

where $\# \mathcal{M}(p_1, p_2, \dots, p_i, q)$ is the cardinality of the moduli space of J -holomorphic disks whose boundary lies in $A_1, A_2, \dots, A_i$ and goes through $p_1, p_2, \dots, p_i, q$ in cyclic order. If the moduli space is not zero dimensional, it is counted as 0.

The above A_∞ category is called the *Fukaya category* of M , and is denoted by $\mathcal{Fuk}(M)$. We shall discuss a little more about it in the last section. However, from the definition, one sees that the A_∞ operators m_i are in general non-trivial, and therefore it is not easy to compute the Hochschild or cyclic homology of $\mathcal{Fuk}(M)$.

Let (M, ω, η, J) be an exact symplectic manifold. A smooth function $f : M \rightarrow \mathbb{C}$ is called a *symplectic Lefschetz fibration* if f have isolated non-degenerate critical points such that near each of these critical points, say p , f has the form

$$f(z_1, z_2, \dots, z_n) = f(p) + z_1^2 + z_2^2 + \dots + z_n^2,$$

and that the fibers of f are symplectic submanifolds of M . Assume that each critical value corresponds to one critical point.

Fix a regular value λ_0 of f , and consider an arc $\gamma \subset \mathbb{C}$ joining λ_0 to a critical value $\lambda_i = f(p_i)$. For the symplectic Lefschetz fibration, there is a natural way of doing symplectic parallel transport along any path, and for γ the set of points transported to p_i forms a Lagrangian disk $\Delta_\gamma \subset M$. The boundary of $\Delta_\gamma|f^{-1}(\lambda_0)$ is therefore a Lagrangian sphere, which is called a *vanishing cycle*. Let $\gamma_1, \gamma_2, \dots, \gamma_r$ be a collection of arcs in $\mathbb{C}$ joining λ_0 to the various critical values of f , and assume they only intersect at λ_0 , and are ordered in the clockwise direction around p_0 . Each arc γ_i gives a Lagrangian sphere $L_i \subset f^{-1}(\lambda_0)$. After a small perturbation we can assume that these spheres intersect each other transversely inside $f^{-1}(\lambda_0)$.

Definition-Lemma 1 (Seidel). Let $f : M \rightarrow \mathbb{C}$ be a symplectic Lefschetz fibration. The directed A_∞ category of vanishing cycles, denoted by $\text{Lag}_{\text{vc}}^\rightarrow(f, \{\gamma_i\})$, is the A_∞ category whose objects are $L_1, L_2, \dots, L_r$ corresponding to the vanishing cycles, and whose morphisms are given by

$$\text{Hom}(L_i, L_j) = \begin{cases} CF^*(L_i, L_j), & \text{if } i < j, \\ k \cdot id, & \text{if } i = j, \\ 0, & \text{if } i > j, \end{cases} \quad (5)$$

with $m_k : \text{Hom}(L_{i_0}, L_{i_1}) \otimes \dots \otimes \text{Hom}(L_{i_{k-1}}, L_{i_k}) \rightarrow \text{Hom}(L_{i_0}, L_{i_k})$ defined by

$$m_k = \begin{cases} m_k|_{\mathcal{Fuk}(M)}, & \text{if } i_0 < i_1 < \dots < i_k, \\ 0, & \text{otherwise.} \end{cases}$$

According to Seidel, any A_∞ category $\mathcal{A}$ whose morphisms between two objects are in the form (5) with $CF^*(-, -)$ replaced by a finite dimensional vector space is called *directed*. Similarly to Lemma 2 we have the following:

Lemma 3. Suppose $\mathcal{A}$ is a directed A_∞ category with n objects, then

$$CH_*(\mathcal{A}) \cong CH_*(k)^{\oplus n}.$$

Corollary 2. *The cyclic homology of the directed Fukaya category $\text{Lag}_{\text{vc}}^\rightarrow(f, \{\gamma_i\})$ of vanishing cycles is isomorphic as vector spaces to the direct sum of r copies of $CH_*(k)$:*

$$CH_*(\text{Lag}_{\text{vc}}^\rightarrow(f, \{\gamma_i\})) \cong CH_*(k)^{\oplus r},$$

where r is the number of critical points.

Seidel's construction of $\text{Lag}_{\text{vc}}^\rightarrow(f, \{\gamma_i\})$ depends on the order of the paths connecting the critical points (hence the order of the vanishing cycles). A key theorem that Seidel proves is that if we change the order of the vanishing cycles, the two sets are related by a sequence of *mutations*. A classical result of triangulated categories says that if two exceptional collections are related by mutations, they generate the the same triangulated category. Thus if we consider the derived A_∞ category $\mathcal{D}(\text{Lag}_{\text{vc}}^\rightarrow(f, \{\gamma_i\}))$ (for a definition, see [32]), where the vanishing cycles form a full strong exceptional collection, then it is independent of the order, and hence is an invariant of the Lefschetz fibration. The cyclic homology of $\text{Lag}_{\text{vc}}^\rightarrow(f, \{\gamma_i\})$, which is isomorphic to $CH_*(k)^{\oplus n}$ as a linear space, is independent of the order of the vanishing cycles, too: a theorem of Keller ([24]) says that the cyclic homology is invariant under derived equivalence.

3. CALABI-YAU A_∞ CATEGORIES

Definition 4 (Calabi-Yau A_∞ category). An A_∞ category $\mathcal{A}$ is called *Calabi-Yau* of degree n if there is degree n graded symmetric pairing

$$\text{Hom}(B, A) \otimes \text{Hom}(A, B) \xrightarrow{\langle \cdot, \cdot \rangle} k$$

for each pair of objects A, B , which is non-degenerate and cyclically invariant, where being *cyclically invariant* means the multilinear map

$$(a_0, a_1, \dots, a_n) \mapsto (-1)^{(n-2)|a_0|} \langle a_0, m_n(a_1, \dots, a_n) \rangle,$$

where $a_i \in \text{Hom}(A_i, A_{i+1})$, $i = 0, 1, \dots, n$ (here and in the following we identify A_{n+1} with A_0), is invariant under the cyclic permutation, i.e.

$$(-1)^{(n-2)|a_0|} \langle a_0, m_n(a_1, \dots, a_n) \rangle = (-1)^{n+|a_n|(|a_0|+\dots+|a_{n-1}|)} \cdot (-1)^{(n-2)|a_n|} \langle a_n, m_n(a_0, \dots, a_{n-1}) \rangle.$$

From the definition, one sees that if a Calabi-Yau A_∞ category has one object, then it is a cyclic A_∞ algebra in the sense of Getzler-Kapranov ([18]). The directed Fukaya category $\text{Lag}_{\text{vc}}^\rightarrow(f, \{\gamma_i\})$ of vanishing cycles in a symplectic Lefschetz fibration in the previous section is in general not Calabi-Yau.

3.1. Topological conformal field theory. According to Costello, Calabi-Yau A_∞ categories play an essential role in the understanding of topological conformal field theories. In the following we briefly recall his results.

Suppose S is a Riemann surface with boundary, and denote its boundary by ∂S . Suppose there is an embedding of disjoint union of intervals and circles to ∂S . Call the images of the intervals the *open boundaries* (denoted by $\{O\}$) and the images of the circles the *closed boundaries* (denoted by $\{C\}$). If the orientation of an open or closed boundary is compatible with the one induced from S , then we say such a boundary is *incoming*; otherwise, it is *outgoing*.

Suppose Λ is a set, called the set of D-branes. We label the components of $\partial S \setminus (\{\text{open boundaries}\} \cup \{\text{closed boundaries}\})$ by the elements of Λ , the D-branes. Note for each open boundary O of S , there

are two D-branes associated to it, one is labeled to the component containing the tail of O (denoted by $t(O)$), and the other is labeled to the component containing the head of O (denoted by $s(O)$).

Riemann surfaces with open and closed boundaries and labelings form a category, where the objects are disjoint unions of circles and intervals together with labels, and the morphisms between two objects are the Riemann surfaces connecting them. The composition of two morphisms comes from the glueing of the outgoing boundaries of the first Riemann surface with the incoming boundaries of the second one. Note that when glueing the open boundaries, the corresponding head and tail labelings should be the same. Denote the category by $\mathcal{M}_\Lambda$, which is a symmetric monoidal category under disjoint unions. Define

$$\mathcal{OC}_\Lambda := C_*(\mathcal{M}_\Lambda),$$

which is again a symmetric monoidal category, where $C_*(-)$ is the chain functor. There are two full subcategories of $\mathcal{OC}_\Lambda$, denoted by $\mathcal{O}_\Lambda$ and $\mathcal{C}$ respectively, one is the subcategory whose objects are purely open and the other is the one whose objects are all closed.

Definition 5 (Topological conformal field theory). (1) An *open-closed topological conformal field theory (TCFT)* is a h-split symmetric monoidal functor from the category $\mathcal{OC}_\Lambda$ to the category of chain complexes;

(2) An *open TCFT* is a h-split symmetric monoidal functor from $\mathcal{O}_\Lambda$ to the category of chain complexes; and similarly,

(3) A *closed TCFT* is a h-split symmetric monoidal functor from $\mathcal{C}$ to the category of chain complexes.

In the above definition, by a h-split (abbreviation of homologically split) functor we mean that $F(\alpha) \otimes F(\beta) \rightarrow F(\alpha \otimes \beta)$ is split on the homology $HF(\alpha) \otimes HF(\beta) \xrightarrow{\cong} HF(\alpha \otimes \beta)$. Denote by Kom the category of chain complexes. There are inclusions $\mathcal{O}_\Lambda \xrightarrow{i} \mathcal{OC}_\Lambda \xleftarrow{j} \mathcal{C}$. The following is the main result of Costello ([10]):

Theorem 5 (Costello). (1) *The category of open TCFTs with the set of D-branes Λ is homotopy equivalent to the category of Calabi-Yau A_∞ categories with objects Λ ;*

(2) *For any open TCFT Φ , the homotopy universal functor $\mathbb{L}i_*\Phi : \mathcal{OC} \rightarrow \text{Kom}$ is h-split, and hence defines an open-closed TCFT;*

(3) *The closed states of the open-closed TCFT in (2) is the Hochschild cohomology of the A_∞ category of (1).*

The proof of this theorem is heavily based on Costello's previous work [9] on the dual decomposition of the moduli spaces; the reader may refer to [10] for more details. Before briefly going over the proof of Costello, let us note that for $\mathcal{C}$, there is a free S^1 action on the incoming and outgoing boundaries. The following definition is due to Getzler (see [16]). To distinguish the S^1 invariant functor with the ordinary one as above, we here add an adjective *equivariant*):

Definition 6 (Equivariant closed TCFT). An equivariant closed TCFT is an h-split symmetric monoidal functor from $\mathcal{C}$ to Kom which is invariant with respect to the S^1 action, that is, a functor from $C_*^{S^1}(\mathcal{M}_{\text{closed}})$ to Kom , where $\mathcal{M}_{\text{closed}}$ is the moduli space of Riemann surfaces with closed boundaries.

Theorem 6. *Let $\mathcal{A}$ be a Calabi-Yau A_∞ category, then the cyclic cochain complex $\text{Cycl}^*(\mathcal{A})$ defines an equivariant TCFT.*

Proof. We follow closely Costello's original argument, which goes roughly as follows: Suppose A is a differential graded symmetric monoidal (DGMS) category, one may view a functor $F : A \rightarrow \text{Kom}$ as a left A module. He argues that if in general $A \rightarrow B$ is a quasi-isomorphism of dgsm categories, then the category of modules over A and the category of modules over B modules are homotopy equivalent. With this result, to study the modules over A , say, it is sufficient to study the modules over B which is homotopy equivalent to A . This allows Costello to construct an explicit model of $\mathcal{O}_\Lambda$ and $\mathcal{OC}_\Lambda$. Namely, he constructs a DGSM category which is quasi-isomorphic to the category $\mathcal{O}_\Lambda$ and then constructs a module over this category which is quasi-isomorphic to $\mathcal{OC}_\Lambda$.

More precisely, denote by $\mathcal{N}_{g,h,r,s}$ the moduli space of Riemann surfaces of genus g , with h boundary components, r marked points in the boundary and s marked points in the interior. Here the r marked points will later be viewed as inputs and outputs of open strings, and the s marked points as closed strings. There is a partial compactification of $\mathcal{N}_{g,h,r,s}$ into an orbifold with corners, which parameterizes the Riemann surfaces as above but possibly with nodes on the boundary. Inside the compactification $\overline{\mathcal{N}}_{g,h,r,s}$ is an orbi-cell complex $D_{g,h,r,s}$, which parameterizes the Riemann surfaces glued from discs, with each of its components has *at most* one interior marked point. We have ([9, 10]):

- The cellular chain complex of $D_{g,h,r,0}$ models $\mathcal{O}_\Lambda$. The discs with marked points on its boundary is nothing but Stasheff's associahedra, from which one deduces that an open TCFT is homotopy equivalent to a Calabi-Yau A_∞ category;
- The chain complex of $D_{g,h,r,s}$, $s \geq 0$, is a module (which is almost free) over that of $D_{g,h,r,0}$. If we fatten the s marked points into circles, namely, consider the moduli space $G_{g,h,r,s}$, where $G_{g,h,r,s}$ is the moduli space of Riemann surfaces glued from discs and annuli such that they are of genus g , with $h + s$ boundary components, r marked points in the first h boundary components and one marked points in the each of the last s boundary components (coming from the annuli). Then the chain complex of $G_{g,h,r,s}$ for all g, h, r, s , gives a model for $\mathcal{OC}_\Lambda$.
- For an annulus with r marked points in its outer boundary, say $a_1, a_2, \dots, a_n$ (in cyclic order), with D-brane labels $s(a_i) = \lambda_i, t(a_{i+1}) = \lambda_{i+1}$, where $\lambda_{n+1} = \lambda_1$, and a marked point a in its inner boundary (viewed as the closed string) which is parallel to the arc between a_n and a_1 , if viewed as an element in the module of the chain complex of $D_{g,h,r,0}$, gives a closed string state, which is exactly

$$\text{Hom}(\lambda_1, \lambda_2) \otimes \text{Hom}(\lambda_2, \lambda_3) \otimes \dots \otimes \text{Hom}(\lambda_n, \lambda_1).$$

Such an identity is compatible with the differential.

Basically this is Costello's proof; however, observe that there are S^1 ways to mark the point in the inner boundary of the annulus, so if we modulo the S^1 action, namely, only consider $D_{g,h,r,s}$ but not $G_{g,h,r,s}$ (this is Costello's original construction in [9]), we get the closed state is exactly an element in the cyclic chain complex. Also, we would prefer to view such annulus a map from the open string states to the closed string states, which then means it is a cyclic cochain. This proves the theorem. $\square$

Remark 2. *There are some technical details in the proof of Costello, which we didn't mention, for example, the gradings. However, according to Costello's compactification of the moduli spaces ([9]), the cyclic cohomology is probably more natural in his argument than the Hochschild cohomology.*

4. THE LIE BIALGEBRA STRUCTURE

In this section we construct the DG Lie bialgebra structure on the cyclic cochain complex of a Calabi-Yau A_∞ category $\mathcal{A}$.

First, we continue the discussion of the signs in §2 by encoding them with the non-degenerate pairing. Note that the pairing $\langle \cdot, \cdot \rangle$ on V induces a symplectic pairing on ΣV : $\omega(\bar{\alpha}, \bar{\beta}) := (-1)^{|\bar{\alpha}|} \langle \alpha, \beta \rangle$. The cyclic invariance of

$$a_0 \otimes a_1 \otimes \cdots \otimes a_n \mapsto (-1)^{(n-2)|a_0|} \langle a_0, m_n(a_1, a_2, \dots, a_n) \rangle$$

implies the following commutative diagram:

$$\begin{array}{ccc} \bar{a}_0 \otimes \bar{a}_1 \otimes \cdots \otimes \bar{a}_n & \xrightarrow{\bar{t}} & (-1)^{|\bar{a}_n|(|\bar{a}_0| + \cdots + |\bar{a}_{n-1}|)} \bar{a}_n \otimes \bar{a}_0 \otimes \cdots \otimes \bar{a}_{n-1} \\ \downarrow & & \downarrow \\ \omega(\bar{a}_0, \bar{m}_n(\bar{a}_1, \dots, \bar{a}_n)) & \equiv & (-1)^{|\bar{a}_n|(|\bar{a}_0| + \cdots + |\bar{a}_{n-1}|)} \omega(\bar{a}_n, \bar{m}_n(\bar{a}_0, \dots, \bar{a}_{n-1})). \end{array}$$

In other words, in the induced multilinear function

$$\bar{a}_0 \otimes \bar{a}_1 \otimes \cdots \otimes \bar{a}_n \mapsto \omega(\bar{a}_0, \bar{m}_n(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)),$$

the sign after the cyclic permutation only comes from the Koszul convention, and the factor $(-1)^n$ from the cyclic permutation does not show up.

4.1. Construction of the Lie bialgebra. Suppose $\mathcal{A}$ is a Calabi-Yau A_∞ category.

Definition 7 (Lie bialgebra). Let L be a (possibly graded) k -space. A Lie bialgebra on L is the triple $(L, [\cdot, \cdot], \delta)$ such that

- $(L, [\cdot, \cdot])$ is a Lie algebra;
- (L, δ) is a Lie coalgebra;
- The Lie algebra and coalgebra satisfy the following identity, called the *Drinfeld compatibility*:

$$\delta[a, b] = \sum_{(a)} ((-1)^{|a''||b|} [a', b] \otimes a'' + a' \otimes [a'', b]) + \sum_{(b)} ([a, b'] \otimes b'' + (-1)^{|a||b'|} b' \otimes [a, b'']),$$

for all $a, b \in L$, where we write $\delta(a) = \sum_{(a)} a' \otimes a''$ and $\delta(b) = \sum_{(b)} b' \otimes b''$.

If moreover, $[\cdot, \cdot] \circ \delta(a) \equiv 0$, for all $a \in L$, $(L, [\cdot, \cdot], \delta)$ is said to be *involutive*. If the Lie bracket has degree k and the Lie cobracket has degree l , denote the Lie bialgebra with degree (l, k) .

Definition-Lemma 2 (Lie algebra). Suppose $\mathcal{A}$ is a Calabi-Yau category of degree n . Define $[\cdot, \cdot] : \text{Cycl}^*(\mathcal{A}) \otimes \text{Cycl}^*(\mathcal{A}) \rightarrow \text{Cycl}^*(\mathcal{A})$ by

$$\begin{aligned} [f, g](\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) &:= \sum_{i < j} \sum_{p, q=1}^{\dim \text{Hom}(A_i, A_{i+1})} \pm f(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{e}_p, \bar{a}_{j+1}, \dots, \bar{a}_n) \cdot g(\omega^{pq} \bar{f}_q, \bar{a}_i, \dots, \bar{a}_j) \\ &\quad - \sum_{i < j} \sum_{p, q=1}^{\dim \text{Hom}(A_i, A_{i+1})} \pm g(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{e}_p, \bar{a}_{j+1}, \dots, \bar{a}_n) \cdot f(\omega^{pq} \bar{f}_q, \bar{a}_i, \dots, \bar{a}_j), \end{aligned}$$

where $\{\bar{e}_p\}$ and $\{\bar{f}_q\}$ are bases of $\Sigma\text{Hom}(A_i, A_{j+1})$ and $\Sigma\text{Hom}(A_{j+1}, A_i)$ respectively, and $(\omega^{pq}) = ((-1)^{|e_p|+1}\langle e_p, f_q \rangle)^{-1}$. Then $(\text{Cycl}^*(\mathcal{A}), [,], b)$ forms a differential graded Lie algebra of degree $2 - n$.

Proof. Recall the cyclic operator $\bar{t}: (\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \mapsto (\bar{a}_n, \bar{a}_1, \dots, \bar{a}_{n-1})$; if we set $N := 1 + t + \dots + t^n$, then $(1 - t)N = N(1 - t) = 0$, and we may write any homogeneous element in $\text{Cycl}^*(\mathcal{A})$ in the form $\alpha = N(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)$, where $\bar{a}_i \in s(\text{Hom}(A_i, A_{i+1})^*)$. The bracket $[,]$ can be written by the following formula: for $\alpha = N(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)$ and $\beta = N(\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m)$,

$$[\alpha, \beta] = \sum_{i=1}^n \sum_{j=1}^m \pm \omega(\bar{a}_i, \bar{b}_j) N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_m, \bar{b}_1, \dots, \bar{b}_{j-1}, \bar{a}_{i+1}, \dots, \bar{a}_n),$$

where ω is the induced symplectic form on $s(\text{Hom}(-, -)^*)$. From the symplecticity of ω , one sees that $[,]$ is graded skew symmetric.

We show the Jacobi identity: for $\alpha = N(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)$, $\beta = N(\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m)$, $\gamma = N(\bar{c}_1, \bar{c}_2, \dots, \bar{c}_p)$ in $\text{Cycl}^*(\mathcal{A})$,

$$\begin{aligned} & [[\alpha, \beta], \gamma] \\ &= \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{a}_k, \bar{c}_l) N(\bar{a}_1, \dots, \bar{b}_{j+1}, \dots, \bar{b}_{j-1}, \dots, \bar{c}_{l+1}, \dots, \bar{c}_{l-1}, \dots, \bar{a}_n) \end{aligned} \quad (6)$$

$$+ \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{b}_k, \bar{c}_l) N(\bar{a}_1, \dots, \bar{b}_{j+1}, \dots, \bar{c}_{l+1}, \dots, \bar{c}_{l-1}, \dots, \bar{b}_{j-1}, \dots, \bar{a}_n). \quad (7)$$

Similarly, we have

$$\begin{aligned} & [[\beta, \gamma], \alpha] \\ &= \sum_{i,j,k,l} \pm \omega(\bar{b}_j, \bar{c}_l) \omega(\bar{b}_k, \bar{a}_i) N(\bar{b}_1, \dots, \bar{c}_{l+1}, \dots, \bar{c}_{l-1}, \dots, \bar{a}_{i+1}, \dots, \bar{a}_{i-1}, \dots, \bar{b}_m) \end{aligned} \quad (8)$$

$$+ \sum_{i,j,k,l} \pm \omega(\bar{b}_j, \bar{c}_l) \omega(\bar{c}_k, \bar{a}_i) N(\bar{b}_1, \dots, \bar{c}_{l+1}, \dots, \bar{a}_{i+1}, \dots, \bar{a}_{i-1}, \dots, \bar{c}_{l-1}, \dots, \bar{b}_m), \quad (9)$$

and

$$\begin{aligned} & [[\alpha, \beta], \gamma] \\ &= \sum_{i,j,k,l} \pm \omega(\bar{c}_l, \bar{a}_i) \omega(\bar{c}_k, \bar{b}_j) N(\bar{c}_1, \dots, \bar{a}_{i+1}, \dots, \bar{a}_{i-1}, \dots, \bar{b}_{j+1}, \dots, \bar{b}_{j-1}, \dots, \bar{c}_p) \end{aligned} \quad (10)$$

$$+ \sum_{i,j,k,l} \pm \omega(\bar{c}_l, \bar{a}_i) \omega(\bar{a}_k, \bar{b}_j) N(\bar{c}_1, \dots, \bar{a}_{i+1}, \dots, \bar{b}_{j+1}, \dots, \bar{b}_{j-1}, \dots, \bar{a}_{i-1}, \dots, \bar{c}_p). \quad (11)$$

By the cyclic invariance of N and symplecticity of ω , (6) cancels with (11), so do (7) with (8) and (9) with (10). This proves the Jacobi identity.

Finally, we show that the bracket commutes with the boundary:

$$\begin{aligned} (b[f, g])(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) &= [f, g](b(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)) \\ &= [f, g]\left(\sum_i \sum_k \pm (\bar{a}_1, \dots, \bar{m}_k(\bar{a}_i, \dots, \bar{a}_{i+k-1}), \dots, \bar{a}_n)\right) \end{aligned} \quad (12)$$

$$+ [f, g]\left(\sum_j \sum_k \pm (\bar{m}_k(\bar{a}_{n-j}, \dots, \bar{a}_n, \bar{a}_1, \dots, \bar{a}_i), \dots, \bar{a}_{n-j-1})\right), \quad (13)$$

while

$$([bf, g] + (-1)^{|f|}[f, bg])(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)$$

$$= \sum_{i,j} \sum_{p,q} \pm f(b(\bar{a}_i, \dots, \bar{a}_j, \bar{e}_p)) \cdot g(\omega^{pq} \bar{e}_q, \bar{a}_{j+1}, \dots, \bar{a}_{i-1}) \quad (14)$$

$$+ \sum_{i,j} \sum_{p,q} \pm f(\bar{a}_i, \dots, \bar{a}_j, \bar{e}_p) \cdot g(b(\omega^{pq} \bar{e}_q, \bar{a}_{j+1}, \dots, \bar{a}_{i-1})). \quad (15)$$

From the definition of $[\cdot, \cdot]$, one sees (14) + (15) contains more terms than (12) + (13), namely, those terms involving $\bar{m}_k$ acting on $\bar{e}_p$. The extra terms coming from (14) are

$$\sum_{p,q} \sum_r f(\bar{a}_{l+1}, \dots, \bar{m}_r(\bar{a}_k, \dots, \bar{a}_j, \bar{e}_p, \bar{a}_i, \dots, \bar{a}_l)) \cdot g(\omega^{pq} \bar{e}_q, \bar{a}_{j+1}, \dots, \bar{a}_{i-1}) \quad (16)$$

and the ones from (15) are

$$\sum_{p,q} \sum_r (\bar{a}_i, \dots, \bar{a}_j, \bar{e}_p) \cdot g(\bar{m}_r(\bar{a}_k, \dots, \bar{a}_{i-1}, \omega^{pq} \bar{e}_q, \bar{a}_{j+1}, \dots, \bar{a}_l), \dots, \bar{a}_{k-1}). \quad (17)$$

However, these two groups of terms cancel with each other: by the non-degeneracy of the pairing,

$$\begin{aligned} & \sum_{p,q} \bar{m}_r(\bar{a}_k, \dots, \bar{a}_j, \bar{e}_p, \bar{a}_i, \dots, \bar{a}_l) \otimes \omega^{pq} \bar{e}_q \\ &= \sum_{p,q} \sum_{s,t} \langle \bar{m}_r(\bar{a}_k, \dots, \bar{a}_j, \bar{e}_p, \bar{a}_i, \dots, \bar{a}_l), \bar{e}_s \rangle \omega^{st} \bar{e}_t \otimes \omega^{pq} \bar{e}_q \\ &= \sum_{p,q} \sum_{s,t} \bar{e}_t \otimes \omega^{st} \cdot \langle \bar{m}_r(\bar{a}_k, \dots, \bar{a}_j, \bar{e}_p, \bar{a}_i, \dots, \bar{a}_l), \bar{e}_s \rangle \omega^{pq} \bar{e}_q \\ &\stackrel{\text{cyclicity}}{=} \sum_{p,q} \sum_{s,t} \pm \bar{e}_t \otimes \omega^{st} \cdot \langle \bar{m}_r(\bar{a}_i, \dots, \bar{a}_l, \bar{e}_s, \bar{a}_k, \dots, \bar{a}_j), \bar{e}_p \rangle \omega^{pq} \bar{e}_q \\ &= \sum_{s,t} \pm \bar{e}_t \otimes \bar{m}_r(\bar{a}_i, \dots, \bar{a}_l, \omega^{st} \bar{e}_s, \bar{a}_k, \dots, \bar{a}_j) \\ &= - \sum_{s,t} \pm \bar{e}_t \otimes \bar{m}_r(\bar{a}_i, \dots, \bar{a}_l, \omega^{ts} \bar{e}_s, \bar{a}_k, \dots, \bar{a}_j). \end{aligned}$$

By substituting the above identity into (16) we get exactly (17), and therefore $(\text{Cycl}^*(\mathcal{A}), [\cdot, \cdot], b)$ is a differential graded Lie algebra. $\square$

Definition-Lemma 3 (Lie coalgebra). Suppose $\mathcal{A}$ is a Calabi-Yau category of degree n . Define $\text{Cycl}^*(\mathcal{A}) \rightarrow \text{Cycl}^*(\mathcal{A}) \otimes \text{Cycl}^*(\mathcal{A})$ by

$$\begin{aligned} & (\delta f)(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \otimes (\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m) \\ &:= \sum_{i=1}^n \sum_{j=1}^m \sum_{p,q=1}^{\dim \text{Hom}(A_i, B_j)} \pm f(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{e}_p, \bar{b}_j, \dots, \bar{b}_m, \bar{b}_1, \dots, \bar{b}_{j-1}, \omega^{pq} \bar{f}_q, \bar{a}_i, \dots, \bar{a}_n), \end{aligned}$$

where $\{\bar{e}_p\}$ and $\{\bar{f}_q\}$ are bases of $\Sigma \text{Hom}(A_i, B_j)$ and $\Sigma \text{Hom}(B_j, A_i)$ respectively, and $(\omega^{pq}) = ((-1)^{|e_p|+1} \langle e_p, f_q \rangle)^{-1}$. Then $(\text{Cycl}^*(\mathcal{A}), \delta, b)$ forms a Lie coalgebra of degree $2 - n$.

Proof. From the definition of δ , the following two statements are obvious:

- (1) δf is well defined, namely, the value of δf is invariant under the cyclic permutations of $(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)$ and $(\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m)$;
- (2) δf is (graded) skew-symmetric, namely, if we switch $(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)$ and $(\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m)$, the sign of the value of δf changes.

If writing $f \in \text{Cycl}^*(V)$ in the form $f = N(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n)$, then

$$\begin{aligned} \delta f &= \delta N(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \\ &= \sum_{i < j} \omega(\bar{a}_i, \bar{a}_j) N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_n) \otimes N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \end{aligned} \quad (18)$$

$$- \sum_{i < j} \omega(\bar{a}_i, \bar{a}_j) N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \otimes N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_n). \quad (19)$$

Therefore

$$\begin{aligned} &(id \otimes \delta) \circ \delta N(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \\ &= \sum_{i < j} \sum_{k < l} \omega(\bar{a}_i, \bar{a}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_n) \\ &\quad \otimes N(\bar{a}_{j+1}, \dots, \bar{a}_{k-1}, \bar{a}_{l+1}, \dots, \bar{a}_{j-1}) \otimes N(\bar{a}_{k+1}, \dots, \bar{a}_{l-1}) \end{aligned} \quad (20)$$

$$\begin{aligned} &- \sum_{i < j} \sum_{k < l} \omega(\bar{a}_i, \bar{a}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_n) \otimes N(\bar{a}_{k+1}, \dots, \bar{a}_{l-1}) \\ &\quad \otimes N(\bar{a}_{j+1}, \dots, \bar{a}_{k-1}, \bar{a}_{l+1}, \dots, \bar{a}_{j-1}) \end{aligned} \quad (21)$$

$$\begin{aligned} &+ \sum_{i < j} \sum_{k < l} \omega(\bar{a}_i, \bar{a}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \otimes N(\bar{a}_1, \dots, \bar{a}_{k-1}, \dots, \bar{a}_{l+1}, \dots, \bar{a}_n) \\ &\quad \otimes N(\bar{a}_{k+1}, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_{l-1}) \end{aligned} \quad (22)$$

$$\begin{aligned} &- \sum_{i < j} \sum_{k < l} \omega(\bar{a}_i, \bar{a}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \otimes N(\bar{a}_{k+1}, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_{l-1}) \\ &\quad \otimes N(\bar{a}_1, \dots, \bar{a}_{k-1}, \dots, \bar{a}_{l+1}, \dots, \bar{a}_n) \end{aligned} \quad (23)$$

$$\begin{aligned} &+ \sum_{i < j} \sum_{k < l} \omega(\bar{a}_i, \bar{a}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \\ &\quad \otimes N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_{k-1}, \bar{a}_{l+1}, \dots, \bar{a}_n) \otimes N(\bar{a}_{k+1}, \dots, \bar{a}_{l-1}) \end{aligned} \quad (24)$$

$$\begin{aligned} &- \sum_{i < j} \sum_{k < l} \omega(\bar{a}_i, \bar{a}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \\ &\quad \otimes N(\bar{a}_{k+1}, \dots, \bar{a}_{l-1}) \otimes N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_{k-1}, \bar{a}_{l+1}, \dots, \bar{a}_n), \end{aligned} \quad (25)$$

where (20) + (21) come from (18) and the other terms from (19).

Let $\tau : \alpha \otimes \beta \otimes \gamma \mapsto \pm \gamma \otimes \alpha \otimes \beta$ be the cyclic permutation of three elements, then

$$\begin{aligned} (20) + \tau^2(22) &= 0, & \tau(20) + (22) &= 0, & \tau^2(20) + \tau(22) &= 0; \\ (21) + \tau(23) &= 0, & \tau(21) + \tau^2(23) &= 0, & \tau^2(21) + (23) &= 0; \\ (24) + \tau^2(25) &= 0, & \tau(20) + (25) &= 0, & \tau^2(24) + \tau(25) &= 0; \end{aligned}$$

that means, $(\tau^2 + \tau + id) \circ (id \otimes \delta) \circ \delta f = 0$. This proves the co-Jacobi identity.

Next, we show that b respects the cobracket. This is similar to the Lie case. By definition,

$$\begin{aligned} &((b \otimes id \pm id \otimes b) \delta(f))((\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \otimes (\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m)) \\ &= \delta f(b(\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \otimes (\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m) \pm (\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \otimes b(\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m)), \end{aligned} \quad (26)$$

while

$$\delta(b(f))((\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n) \otimes (\bar{b}_1, \bar{b}_2, \dots, \bar{b}_m))$$

$$= \sum_{i=1}^n \sum_{j=1}^m \sum_{p,q} \pm b(f)(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{e}_p, \bar{b}_j, \dots, \bar{b}_m, \bar{b}_1, \dots, \bar{b}_{j-1}, \omega^{pq} \bar{e}_q, \bar{a}_i, \dots, \bar{a}_n) \quad (27)$$

$$- \sum_{j=1}^m \sum_{i=1}^n \sum_{p,q} \pm b(f)(\bar{b}_1, \dots, \bar{b}_{j-1}, \bar{e}_p, \bar{a}_i, \dots, \bar{a}_n, \bar{a}_1, \dots, \bar{a}_{i-1}, \omega^{pq} \bar{e}_q, \bar{b}_j, \dots, \bar{b}_m). \quad (28)$$

Compared with (26), (27) + (28) has more terms

$$\sum_{p,q} f(\bar{a}_1, \dots, \bar{m}_r(\bar{a}_k, \dots, \bar{e}_p, \dots, \bar{b}_{l-1}), \bar{b}_l, \dots, \omega^{pq} \bar{e}_q, \dots, \bar{a}_n) \quad (29)$$

$$+ \sum_{p,q} f(\bar{a}_1, \dots, \bar{e}_p, \dots, \bar{b}_{j-1}, \bar{m}_r(\bar{b}_j, \dots, \omega^{pq} \bar{e}_q, \dots, \bar{a}_l), \dots, \bar{a}_n) \quad (30)$$

$$+ \sum_{p,q} f(\bar{a}_1, \dots, \bar{m}_r(\bar{a}_k, \dots, \bar{e}_p, \bar{b}_j, \dots, \omega^{pq} \bar{e}_q, \dots, \bar{a}_{l-1}), \bar{a}_l, \dots, \bar{a}_n), \quad (31)$$

However, a similar argument as in the Lie case, (29) + (30) vanishes, and the terms in (31) come in pair (counting e_p and e_q), which cancel within themselves. This proves that the differential commutes with the cobracket. $\square$

Theorem 7. *Let $\mathcal{A}$ be a Calabi-Yau A_∞ category of degree n . The cyclic cochain complex $\text{Cycl}^*(\mathcal{A})$ has the structure of a DG involutive Lie bialgebra of degree $(2-n, 2-n)$.*

Proof. From the above three definitions, we only need to prove that $(\text{Cycl}^*(\mathcal{A}), [\cdot, \cdot], \delta)$ satisfies the Drinfeld compatibility and the involutivity. Let $\alpha = N(\bar{a}_1, \dots, \bar{a}_n)$ and $\beta = N(\bar{b}_1, \dots, \bar{b}_m)$, and write $\delta(\alpha) = \alpha^{(1)} \otimes \alpha^{(2)}$ and $\delta(\beta) = \beta^{(1)} \otimes \beta^{(2)}$.

We have

$$[\alpha, \beta] = \sum_{i,j} \pm \omega(\bar{a}_i, \bar{b}_j) N(\bar{a}_{i+1}, \dots, \bar{a}_i, \bar{a}_1, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_m, \bar{b}_1, \dots, \bar{b}_{j-1})$$

and

$$\begin{aligned} & \delta[\alpha, \beta] \\ &= \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_{k+1}, \dots, \bar{a}_{l-1}) \otimes N(\bar{a}_{l+1}, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_{j-1}, \bar{a}_{i+1}, \dots, \bar{a}_{k-1}) \\ &+ \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_{k+1}, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_{l-1}) \otimes N(\bar{b}_{l+1}, \dots, \bar{b}_{j-1}, \bar{a}_{i+1}, \dots, \bar{a}_{k-1}) \\ &+ \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{a}_{k+1}, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_{j-1}, \bar{a}_{i+1}, \dots, \bar{a}_{l-1}) \otimes N(\bar{a}_{l+1}, \dots, \bar{a}_{k-1}) \\ &+ \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{b}_{k+1}, \dots, \bar{b}_{l-1}) \otimes N(\bar{b}_{l+1}, \dots, \bar{b}_{j-1}, \bar{a}_{i+1}, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_{k-1}) \\ &+ \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{b}_{k+1}, \dots, \bar{b}_{j-1}, \bar{a}_{i+1}, \dots, \bar{a}_{l-1}) \otimes N(\bar{a}_{l+1}, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_{k-1}) \\ &+ \sum_{i,j,k,l} \pm \omega(\bar{a}_i, \bar{b}_j) \omega(\bar{a}_k, \bar{a}_l) N(\bar{b}_{k+1}, \dots, \bar{b}_{j-1}, \bar{a}_{i+1}, \dots, \bar{a}_{i-1}, \bar{b}_{j+1}, \dots, \bar{b}_{l-1}) \otimes N(\bar{b}_{l+1}, \dots, \bar{b}_{k-1}). \end{aligned}$$

In the above, the second summation and the fifth summation cancel with each other. The first summation is equal to $\alpha^{(1)} \otimes [\alpha^{(2)}, \beta]$; the third summation is equal to $[\alpha^{(1)}, \beta] \otimes \alpha^{(2)}$; the fourth summation

is equal to $\beta^{(1)} \otimes [\alpha, \beta^{(2)}]$; and the sixth summation is equal to $[\alpha, \beta^{(1)}] \otimes \beta^{(2)}$. Thus we obtain the Drinfeld compatibility.

Let $\alpha = N(\bar{a}_1, \dots, \bar{a}_n)$, then

$$\begin{aligned} \delta(\alpha) &= \sum_{i < j} \pm \omega(\bar{a}_i, \bar{a}_j) N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_n) \otimes N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \\ &\quad - \sum_{i < j} \pm \omega(\bar{a}_i, \bar{a}_j) N(\bar{a}_{i+1}, \dots, \bar{a}_{j-1}) \otimes N(\bar{a}_1, \dots, \bar{a}_{i-1}, \bar{a}_{j+1}, \dots, \bar{a}_n). \end{aligned}$$

Noting that the two terms in the right hand side are anti-symmetric and by a similar argument as above, one obtains that $[\cdot, \cdot] \circ \delta = 0$ holds identically. $\square$

Geometrically, for a Calabi-Yau A_∞ category $\mathcal{A}$, if we view $\mathcal{A}$ as an open TCFT, then the bracket and cobracket are given by the gluing and splitting of the open states of $\mathcal{A}$ in all possible ways. The construction is inspired from string topology (see [5] and also §4.3), and some of the above computations are similar to [7].

4.2. Noncommutative symplectic geometry. Our construction of the Lie bialgebra was inspired by string topology (see [5, 36] and §4.3). After the first version of the paper was put on arXiv, we were informed by Kontsevich that he and Soibelman ([27, §10]) have a nice cohomological characterization of the cyclic structure, and our formula can also be deduced from theirs. Some related results are Kontsevich [26], Ginzburg [19], Schedler [35] and Hamilton [20]; an essentially equivalent result also appears in Barannikov [3]. In this subsection we would like to study this relationship in more detail.

Definition 8 (Noncommutative differential and de Rham forms). Let V be a vector space. The module of noncommutative 1-forms $\Omega^1(V)$ is defined by

$$\Omega^1(V) := T(V^*) \oplus T^+(V^*).$$

There is a $T(V^*)$ -bimodule structure on $\Omega^1(V)$ which is given by

$$a \cdot (x \otimes y) := ax \otimes y; \quad (x \otimes y) \cdot a := x \otimes ya - xy \otimes a,$$

for $a, x \in T(V^*)$ and $y \in T^+(V^*)$. Define $d : T(V^*) \rightarrow \Omega^1(V)$ by

$$d(x) := 1 \otimes x, \quad d(1) = 0.$$

Let $A := T(V^*)$. The algebra of *noncommutative differential forms* $\Omega^\bullet(V)$ are defined by

$$\Omega^\bullet(V) := T_A(\Sigma\Omega^1(V)) = A \oplus \bigoplus_{n=1}^{\infty} \underbrace{\Sigma\Omega^1(V) \otimes_A \cdots \otimes_A \Sigma\Omega^1(V)}_n.$$

Lifting d to $\Omega^\bullet(V)$ gives a differential graded associative algebra structure on $\Omega^\bullet(V)$. The *noncommutative de Rham forms* $DR^\bullet(V)$ are defined by

$$DR^\bullet(V) := \frac{\Omega^\bullet(V)}{[\Omega^\bullet(V), \Omega^\bullet(V)]},$$

with the induced differential, still denoted by d .

Definition 9 (Lie derivative and contraction). Let V be a vector space and $\Omega^\bullet(V)$ be as above, and let $\xi : T(V^*) \rightarrow T(V^*)$ be a vector field.

(1) The *Lie derivative* $L_\xi : \Omega^\bullet(V) \rightarrow \Omega^\bullet(V)$ with respect to ξ is defined by:

$$L_\xi(x) := \xi(x), \quad L_\xi(dx) := (-1)^{|\xi|} d(\xi(x)), \quad \text{for } x \in T(V^*).$$

(2) The *contraction* $\iota_\xi : \Omega^\bullet(V) \rightarrow \Omega^\bullet(V)$ of ξ is defined by:

$$\iota_\xi(x) := 0, \quad \iota_\xi(dx) := \xi(x), \quad \text{for } x \in T(V^*).$$

The Lie derivative and contraction preserve the commutators and hence descend to $DR^\bullet(V)$.

Definition 10 (Symplectic form and symplectic field). Let V be a vector space. A 2-form $\omega \in DR^2(V)$ is called *symplectic* if

- (1) ω is closed, i.e. $d\omega = 0$;
- (2) ω is non-degenerate, i.e. there is a bijection:

$$\begin{aligned} Der(T(V^*)) &\longrightarrow DR^1(V), \\ \xi &\longmapsto \iota_\xi(\omega). \end{aligned}$$

A vector field $\xi : T(V^*) \rightarrow T(V^*)$ is called a *symplectic vector field* if $L_\xi(\omega) = 0$.

A symplectic form $\omega \in DR^2(V)$ is called a *constant form* if it can be written as $\sum_i dx_i \wedge dy_i$ for some functions $x_i, y_i \in V^*$. The following theorem is due to Kontsevich and others cited above:

Theorem 8 (Lie bialgebra on the noncommutative 0-forms). *Let V be a vector space with a constant symplectic form ω . The symplectic form on V induces a symplectic form on V^* via the identity $V \rightarrow V^* : v \mapsto \omega(v, \cdot)$, which is still denoted by ω . Define $\{, \} : DR^0(V) \otimes DR^0(V) \rightarrow DR^0(V)$ and $\Delta : DR^0(V) \rightarrow DR^0(V) \otimes DR^0(V)$ as follows: for $a_1, a_2, \dots, a_n; b_1, b_2, \dots, b_m \in V^*$,*

$$\{[a_1 a_2 \cdots a_n], [b_1 b_2 \cdots b_m]\} = \sum_i \sum_j \pm \omega(a_i, b_j) ([a_1 \cdots a_{i-1} b_{j+1} \cdots b_m b_1 \cdots b_{j-1} a_{i+1} \cdots a_n]),$$

and

$$\begin{aligned} &\Delta([a_1 a_2 \cdots a_n]) \\ &= \frac{1}{2} \sum_{i < j} \pm \omega(a_i, a_j) ([a_1 \cdots a_{i-1} a_{j+1} \cdots a_n]) \otimes ([a_{i+1} \cdots a_{j-1}]) \\ &+ \frac{1}{2} \sum_{i < j} \pm \omega(a_i, a_j) ([a_{i+1} \cdots a_{j-1}]) \otimes ([a_1 \cdots a_{i-1} a_{j+1} \cdots a_n]). \end{aligned}$$

Then $(DR^0(V), \{, \}, \Delta)$ forms an involutive Lie bialgebra.

In terms of the terminologies introduced above, the Lie bracket is identical to the Poisson bracket on the noncommutative 0-forms. The reader may refer to the above listed references for more details.

So far in this subsection we have assumed only that V is a vector space. Now suppose V is a cyclic A_∞ space. Let $g : V \otimes V$ be the non-degenerate pairing on V , which induces a symplectic form ω on ΣV . There is an isomorphism (Quillen [31, Lemma 1.2, pp. 144])

$$\text{Cycl}^*(V) = \ker(1 - T) \cong \text{coker}(1 - T),$$

namely, the cyclic cochain complex of V is isomorphic to the quotient commutator space. With this identity, the Lie bialgebra defined in the cyclic cochain complex is exactly the one for $(\Sigma V, \omega)$ defined above. Thus we obtain:

Proposition 1. *Let $(V, \{m_i\}, g)$ be a cyclic A_∞ algebra over k , and denote by ω the induced symplectic form on ΣV . Then cyclic cochain complex $\text{Cycl}^*(V)$ is isomorphic to the noncommutative 0-forms on ΣV as Lie bialgebras, where the Lie bialgebra of the former is given in Theorem 7 and the Lie bialgebra of the latter is given in Theorem 8.*

4.3. Example from string topology. Let M be a compact, smooth manifold and LM its free loop space. In [5] (see also [36]) Chas and Sullivan showed, under the program of string topology, that the equivariant homology of the free loop space LM has an involutive Lie bialgebra structure. Such a Lie bialgebra even holds on the chain level, which can be roughly described as follows: Suppose α, β are two transversal equivariant chains in LM (that means, they are in general position); one tries to move the base points of α and β along themselves. At time t_1 for α and t_2 for β , one forms a new chain whose base points are the intersections of these two base points at t_1 and t_2 , and whose fiber is the concatenation of the loops from α and β . By the S^1 invariance of α and β , this new chain is S^1 invariant, too. The Lie bracket of α and β (called the *string bracket*) is thus defined by considering all t_1 and t_2 in S^1 . The Lie cobracket of an equivariant chain, say α , is defined as follows: again, assume α is in general position, one moves the base points of α along itself. Suppose at time t_1 and t_2 , the base points intersect, then one forms two new chains in LM whose base points are the intersections, and whose fibers are the arcs in α from t_1 to t_2 and the rest (note that both arcs are now circles), respectively. Chas and Sullivan argued that with the Lie bracket and cobracket defined this way, the equivariant chain complex of LM forms an involutive Lie bialgebra over $\mathbb{Z}$.

There are some difficulties in the understanding of the Lie bialgebra of string topology, especially in its definition the partially defined transversal intersection is used. However, we may still get some idea from the constructions in the previous sections. First, with the work of K.-T. Chen ([6]) and J. D. S. Jones ([22]), it is nowadays known to topologists that:

Theorem 9 (Chen; Jones). *Suppose M is a simply connected manifold and $C^*(M)$ is a cochain model of M , then the Hochschild (resp. cyclic) chain complex of $C^*(M)$ is quasi-isomorphic to the ordinary (resp. equivariant) chain complex of LM .*

As a corollary, the cyclic cochain complex of $C^*(M)$ is then quasi-isomorphic to the equivariant chain complex of LM .

Recently, Hamilton and Lazarev constructed in [21, Section 5] a finite dimensional cyclic C_∞ algebra over $\mathbb{Q}$, modeling the cochain complex of M . A cyclic C_∞ algebra is a homotopy commutative associative algebra with an invariant bilinear form, it is automatically a cyclic A_∞ algebra.

Theorem 10. *Let M be a simply connected manifold and V be the cyclic A_∞ algebra constructed by Hamilton and Lazarev that models the cochain complex of M . Then the Lie bialgebra of V given in Theorem 1 models the one of string topology tensored with the rational numbers.*

Proof. First observe that the Hochschild and cyclic (co)homology groups are invariant under quasi-isomorphic A_∞ algebras (see for example [17, Lemma 5.3]). There is an isomorphism on the cyclic cohomology groups of $C^*(M)$ and the A_∞ algebra V of Hamilton and Lazarev.

The operator $N : T(sV^*) \rightarrow T(sV^*)$ exactly models the base points rotation of Chas and Sullivan, and the Lie bracket and cobracket given in Section 3 then models the concatenation and splitting of loops over the intersection points. $\square$

Remark 3. *A similar construction of the Lie bialgebra using the cochain model of Lambrechts and Stanley ([28]) which is a strict DG commutative algebra with a non-degenerate pairing has been obtained by the author with Eshmatov and Gan ([7]).*

5. APPLICATION IN SYMPLECTIC TOPOLOGY

In this section we give an example of the Lie bialgebra of Calabi-Yau A_∞ categories that arise from symplectic topology. The details are quite involved, so in the following we will only sketch the main idea.

We start with the definition of the Maslov class of Lagrangian submanifolds. Let $(\mathbb{R}^{2n}, \omega)$ be the standard symplectic vector space. Denote by $\text{Lag}_n = \text{Lag}(\mathbb{R}^{2n}, \omega)$ the set of all linear Lagrangian subspaces. It is known that $\pi_1(\text{Lag}_n) \cong \mathbb{Z}$.

Suppose M is an exact symplectic manifold, and L is an exact Lagrangian submanifold (see Example 2 in §2.1). Let $\phi : (D^2, \partial D^2) \rightarrow (M, L)$ be a map representing $\pi_2(M, L)$. For each $t \in \partial D^2$ we have a Lagrangian subspace $T_{\phi(t)}L \subset T_{\phi(t)}M$, which gives a map $S^1 \rightarrow \text{Lag}_n$. It determines an element in $\pi_1(\text{Lag}_n) \cong \mathbb{Z}$, which is called the *Maslov index* of $[\phi]$, and is denoted by $\mu([\phi])$. The map $[\phi] \mapsto \mu([\phi])$ is called the *Maslov class* of L .

In the following we assume all the exact Lagrangian submanifolds have vanishing Maslov classes. It is proved by symplectic topologists (see, for example, [13, Theorem 1.1] and [32, §8]) that morphism spaces of the Fukaya category of Lagrangian submanifolds are vector spaces over $k = \mathbb{Q}$ or $\mathbb{R}$, and are $\mathbb{Z}$ graded. Moreover, there is a pairing of degree n on the morphism spaces

$$\text{Hom}(A, B) \otimes \text{Hom}(B, A) \rightarrow \text{Hom}(A, A) \xrightarrow{\text{tr}} k$$

induced from the Poincaré duality, making the Fukaya category into a Calabi-Yau A_∞ category. As a corollary to Theorem 7, we have the following:

Theorem 11. *Let M be an exact symplectic $2n$ -manifold with contact type boundary. The cyclic cohomology of the Fukaya category $\mathcal{Fuk}(M)$ of exact Lagrangian submanifolds has the structure of an involutive Lie bialgebra.*

5.1. Symplectic field theory. Symplectic field theory is a program initiated by Eliashberg, Givental and Hofer more than ten years ago ([11]). It has been widely studied by mathematicians and has obtained many interesting results in symplectic and contact topology. In the following we will touch only a small part which is necessary for our discussion.

Let (W, η) be a co-oriented contact manifold, where η is the contact form. There are two concepts associated to W :

- The *symplectization* of W is, by definition, $W \times \mathbb{R}$ with symplectic form $d(e^s \eta)$, where s is the coordinate of the factor $\mathbb{R}$.
- A *Reeb orbit* in W is a closed orbit of the Reeb vector field Y :

$$\eta(Y) = 1, \quad \iota(Y)d\eta = 0.$$

If the contact form η is generic, then the set of Reeb orbits is discrete. To each Reeb orbit, one may assign an integer, called the *Conley-Zehnder index* of the orbit, which is a version of Maslov index ([11, §1.2]).

Choose a generic contact form and consider a J -holomorphic curve $\phi : \Sigma \rightarrow W \times \mathbb{R}$. A theorem of Hofer says that if ϕ has non-removable singular points, then these singular points can only approach the Reeb orbits in W at $\pm\infty$. Moreover, since the Reeb orbits are discrete, one may consider the moduli space of J -holomorphic curves with a given type. Namely, let $\mathcal{M}(\Sigma; q_1^+, q_2^+, \dots, q_m^+; q_1^-, q_2^-, \dots, q_n^-)$ be the set of J -holomorphic curves $\Sigma \rightarrow W \times \mathbb{R}$ such that the only singular points of Σ are $q_1^+, q_2^+, \dots, q_m^+$ at $+\infty$ and $q_1^-, q_2^-, \dots, q_n^-$ at $-\infty$. It is proved in symplectic field theory that the union of $\mathcal{M}(\dots)$'s with all singular type forms a stratified space, whose codimension great than zero strata of a given singular type is described by the “broken” curves, which, topologically, can be realized as J -holomorphic curves which are stretched to be infinitely long in the middle of $W \times \mathbb{R}$ at some time s . $\mathcal{M}(\dots)$ always admits an $\mathbb{R}$ action, which shifts the J -holomorphic curves along the $\mathbb{R}$ factor in $W \times \mathbb{R}$.

Now suppose M is a symplectic manifold with contact type boundary W . The *linearized contact homology* of W is defined as follows (for more details see [8]): let $\text{Cont}_*^{\text{lin}}(W)$ be the linear space spanned by the Reeb orbits (technically, they should be *good* in the sense of [11]), graded by the Conley-Zehnder indices. Define a linear operator

$$d : \text{Cont}_*^{\text{lin}}(W) \rightarrow \text{Cont}_{*-1}^{\text{lin}}(W)$$

by

$$d(q^+) = \sum_{q^- : |q^-| = |q^+| - 1} \#(\mathcal{M}(\text{cylinder}; q^+; q^-)/\mathbb{R}) \cdot q^-.$$

The compactness of $\mathcal{M}(\dots)$ makes d into a differential. The following is achieved in symplectic field theory by Cieliebak and Latschev ([8, Theorem A]):

Theorem 12 (Cieliebak and Latschev). *Let M be a symplectic manifold with contact type boundary W . The chain complex $\text{Cont}_*^{\text{lin}}(W)$ has the structure of a bi-Lie $_\infty$ algebra, which, induces on the linearized contact homology of W an involutive Lie bialgebra of degree $(-1, 5 - 2n)$.*

A bi-Lie $_\infty$ algebra is a homotopy version of an involutive Lie bialgebra, where the Jacobi identity, co-Jacobi identity, Drinfeld compatibility and involutivity all hold up to homotopy (we shall not go to the details of bi-Lie $_\infty$ algebras; for some more discussions, see [8]). Basically, the Lie bracket is given by counting the J -holomorphic spheres with two non-removable singular points at $+\infty$ and one non-removable singular point at $-\infty$ in $W \times \mathbb{R}$. Namely, suppose q_1, q_2 are two Reeb orbits in W . Then the Lie bracket

$$[q_1, q_2] := \sum_q \#(\mathcal{M}(\text{sphere} \setminus \{3 \text{ points}\}; q_1^+, q_2^+; q^-)/\mathbb{R}) \cdot q,$$

where $\mathcal{M}(\text{sphere} \setminus \{3 \text{ points}\}; q_1^+, q_2^+; q^-)$ is the moduli space of J -holomorphic spheres with 3 non-removable singular points, two of which approach q_1 and q_2 in W at $+\infty$ and the third one approaches q in W at $-\infty$. By considering the J -holomorphic spheres with 3 non-removable singular points at $+\infty$ and 1 non-removable singular point at $-\infty$ in W , one obtains that the Jacobi identity for the Lie bracket holds up to homotopy.

In a similar fashion one may define the Lie cobracket and prove that the co-Jacobi identity holds up to homotopy, too. The involutivity and Drinfeld compatibility hold up to homotopy due to the study of the moduli space of J -holomorphic tori with 2 non-removable singular points and J -holomorphic spheres with 4 non-removable singular points (2 at $+\infty$ and 2 at $-\infty$) respectively.

In the theorem of Cieliebak-Latschev, if we shift the degrees of the Reeb orbits up by $n - 3$, then the Lie bialgebra on the linearized contact homology has degree $(2 - n, 2 - n)$. In the following we take this degree shifting.

5.2. From Fukaya category to symplectic field theory. Let M be an exact symplectic manifold with contact type boundary W . M has a *symplectic completion*, which is

$$M \cup_{id: W \rightarrow W \times \{0\}} (W \times \mathbb{R}^{\geq 0}),$$

and is denoted by $\tilde{M}$. $\tilde{M}$ is a symplectic manifold whose symplectic form is induced from M and $W \times \mathbb{R}^{\geq 0}$.

Theorem 13. *Let M be a symplectic manifold with contact type boundary W , and let $\tilde{M}$ be the symplectic completion. Let*

$$f : \text{Cont}_*^{\text{lin}}(W) \rightarrow \text{Cycl}^*(\mathcal{Fuk}(M))$$

be the map such that the value of $f(q)$, $q \in \text{Cont}_(W)$, at $(a_1, a_2, \dots, a_n) \in \text{Hom}(A_1, A_2) \otimes \text{Hom}(A_2, A_3) \otimes \dots \otimes \text{Hom}(A_n, A_1)$ is the cardinality of J -holomorphic disks whose boundary lies in $A_1 \cup A_2 \cup \dots \cup A_n$ and goes through $a_1, a_2, \dots, a_n$ in cyclic order and whose non-removable singular point in $\tilde{M}$ at $+\infty$ approaches q . Then f is a map of chain complexes and induces on the homology a map of Lie bialgebras.*

Sketch of proof. First, we show f is a chain map. For any $q \in \text{Cont}_*(W)$, and $(a_1, a_2, \dots, a_n) \in \text{Hom}(A_1, A_2) \otimes \text{Hom}(A_2, A_3) \otimes \dots \otimes \text{Hom}(A_n, A_1)$, consider the moduli space of J -holomorphic disks with boundaries in $A_1 \cup A_2 \cup \dots \cup A_n$ and through $a_1, a_2, \dots, a_n$ in cyclic order and with a non-removable singular point approaching q . Suppose the moduli space is 1-dimensional, then its boundary is composed of the following two types of broken J -holomorphic curves:

- The first type is a broken curve whose first piece is a J -holomorphic cylinder in $W \times \mathbb{R}$ with singular points approaching q at $+\mathbb{R}$ and q' at $-\infty$, and whose second piece is a J -holomorphic disk with its boundary lying in $A_1 \cup A_2 \cup \dots \cup A_n$ as before and with a non-removable singular point approaching q' in $\tilde{M}$. In other words, it comes from stretching the original singular J -holomorphic disk in $\mathbb{R}^+$ direction at some middle point in $\tilde{M}$;
- The second type is a broken curve whose first piece is a J -holomorphic disk with its boundary lying in $A_1 \cup \dots \cup A_i \cup A_j \cup \dots \cup A_n$ and going through $a_1, \dots, a_{i-1}, p, a_j, \dots, a_n$ in cyclic order and with a non-removable singular point approaching q at $+\infty$, and whose second piece is a J -holomorphic curve with its boundary lying in $A_i \cup A_{i+1} \cup \dots \cup A_j$ and going through $p, a_i, a_{i+1}, \dots, a_{j-1}$, where $p \in A_i \cap A_j$. In other words, it comes from the bubbling-off (of disks) from the original singular J -holomorphic disk.

Now to show $f(dq) = b(f(q))$, consider their values at $(a_1, a_2, \dots, a_n) \in \text{Hom}(A_1, A_2) \otimes \text{Hom}(A_2, A_3) \otimes \dots \otimes \text{Hom}(A_n, A_1)$. Note $f(dq)(a_1, a_2, \dots, a_n)$ is exactly the number of broken J -holomorphic curves of the first type and $b(f(q))(a_1, a_2, \dots, a_n) = f(q)(b(a_1, a_2, \dots, a_n))$ is exactly the number of broken J -holomorphic curves of the second type. It follows that their cardinalities are equal.

Next we show that f maps the bracket to the bracket up to homotopy, which induces a Lie algebra map on the homology. For $q_1, q_2 \in \text{Cont}_*^{\text{lin}}(W)$ and $(a_1, a_2, \dots, a_n) \in \text{Hom}(A_1, A_2) \otimes \text{Hom}(A_2, A_3) \otimes \dots \otimes \text{Hom}(A_n, A_1)$, we now consider J -holomorphic disks with their boundary in $A_1 \cup A_2 \cup \dots \cup A_n$ and through $a_1, a_2, \dots, a_n$ in cyclic order, and having two non-removable singular

points approaching q_1, q_2 in W at $+\infty$. Suppose the moduli space is 1-dimensional, then its boundary points are composed of four types of broken J -holomorphic curves (imaging a pair of pants are placed vertically with two sleeves at $+\infty$ and one at 0):

- The broken point of the pants occurs at one of the upper sleeves but not the other; in other words, one of them is stretched to be infinitely long.
- The broken point of the pants occurs at the bottom sleeve; in other words, there is a bubbling off of a disk at the bottom sleeve.
- The broken point of the pants occurs at somewhere between the saddle point and the bottom sleeve;
- The broken point occurs when the saddle point is pushed down to 0.

These four cases corresponds to the following operations respectively: (1) $f_2(dq_1, q_2)$ and $f_2(q_1, dq_2)$; (2) $b(f_2[q_1, q_2])$; (3) $f[q_1, q_2]$; and (4) $[f(q_1), f(q_2)]$. Here by $f_2(-, -)$ we mean counting the J -holomorphic disks with two non-removable singular points. It follows then that

$$f[q_1, q_2] - [f(q_1), f(q_2)] = f_2(dq_1, q_2) + f_2(q_1, dq_2) - bf_2(q_1, q_2),$$

that is, f is a Lie algebra map up to homotopy.

Finally we show that f maps the cobracket to the cobracket up to homotopy. By considering J -holomorphic cylinders with both boundaries fixed on two sets of cyclic chain complex in $\mathcal{Fuk}(M)$ and with 1 non-removable singular point approaching a Reeb orbit in W at $+\mathbb{R}$ (or equivalently, the J -holomorphic pants with two sleeves fixed in the Lagrangian submanifolds and one sleeve going to $+\infty$), with the similar argument as above, one sees that f is a Lie coalgebra map up to homotopy. $\square$

From the proof, one sees that if we consider all J -holomorphic curves with several non-removable singular points and several boundaries, then f is in fact a map of bi-Lie $_\infty$ algebras. It is interesting then to ask when f induces an isomorphism of Lie bialgebras on the homology level. We hope to address this problem somewhere else.

REFERENCES

- [1] D. Auroux, L. Katzarkov and D. Orlov, *Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves*. Invent. Math. 166 (2006), no. 3, 537-582.
- [2] D. Auroux, L. Katzarkov and D. Orlov, *Mirror symmetry for weighted projective planes and their noncommutative deformations*. Ann. of Math. (2) 167 (2008), no. 3, 867-943.
- [3] S. Barannikov, *Modular operads and Batalin-Vilkovisky geometry*. Int. Math. Res. Not. IMRN 2007, no. 19, Art. ID rnm075.
- [4] A. I. Bondal and M. M. Kapranov, *Enhanced triangulated categories*. (Russian) Mat. Sb. 181 (1990), no. 5, 669-683; translation in Math. USSR-Sb. 70 (1991), no. 1, 93-107.
- [5] M. Chas and D. Sullivan, *Closed string operators in topology leading to Lie bialgebras and higher string algebra*, in *The legacy of Niels Henrik Abel*, 771-784, Springer, Berlin, 2004.
- [6] K.-T. Chen, *Iterated path integrals*, Bulletin of the AMS, Volume 83, Number 5, September 1977, 831-879.
- [7] X. Chen, F. Eshmatov and W. L. Gan, *Quantization of the Lie bialgebra of string topology*, 2009 (arXiv:0902.2161).
- [8] K. Cieliebak and J. Latschev, *The role of string topology in symplectic field theory*, 2007 (arXiv:0706.3284).
- [9] K. Costello, *A dual version of the ribbon graph decomposition of moduli space*. Geom. Topol. 11 (2007), 1637-1652.
- [10] K. Costello, *Topological conformal field theories and Calabi-Yau categories*. Adv. Math. 210 (2007), no. 1, 165-214.
- [11] Y. Eliashberg, A. Givental and H. Hofer, *Introduction to symplectic field theory*. GAFA 2000 (Tel Aviv, 1999). Geom. Funct. Anal. 2000, Special Volume, Part II, 560-673.

- [12] K. Fukaya, *Floer homology and mirror symmetry. II*, in *Minimal surfaces, geometric analysis and symplectic geometry* (Baltimore, MD, 1999), 31-127, Adv. Stud. Pure Math., 34, Math. Soc. Japan, Tokyo, 2002.
- [13] K. Fukaya, *Cyclic symmetry and adic convergence in Lagrangian Floer theory*, 2009 (arXiv:0907.4219).
- [14] S. I. Gelfand and Y. I. Manin, *Methods of homological algebra*. Second edition. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003.
- [15] E. Getzler, *Batalin-Vilkovisky algebras and two-dimensional topological field theories*. Comm. Math. Phys. 159 (1994), no. 2, 265-285.
- [16] E. Getzler, *Two-dimensional topological gravity and equivariant cohomology*. Comm. Math. Phys. 163 (1994), no. 3, 473-489.
- [17] E. Getzler and J.D.S. Jones, *A_∞ -algebras and the cyclic bar complex*. Illinois J. Math. 34 (1990), no. 2, 256-283.
- [18] E. Getzler and M.M. Kapranov, *Cyclic operads and cyclic homology*, in *Geometry, topology and physics*, International Press, Cambridge, MA, 1995, pp. 167-20.
- [19] V. Ginzburg, *Non-commutative symplectic geometry, quiver varieties, and operads*. Math. Res. Lett. 8 (2001), no. 3, 377-400.
- [20] A. Hamilton, *Noncommutative geometry and compactifications of the moduli space of curves*, 2008, (arXiv:0801.0904).
- [21] A. Hamilton and A. Lazarev, *Homotopy algebras and noncommutative geometry*, 2004 (arxiv:math/0410621v1).
- [22] J.D.S. Jones, *Cyclic homology and equivariant homology*. Invent. Math. 87 (1987), 403-423.
- [23] M. M. Kapranov, *On the derived categories of coherent sheaves on some homogeneous spaces*. Invent. Math. 92 (1988), no. 3, 479-508.
- [24] B. Keller, *Invariance and localization for cyclic homology of DG algebras*. J. Pure Appl. Algebra 123 (1998), no. 1-3, 223-273.
- [25] B. Keller, *On the cyclic homology of ringed spaces and schemes*. Doc. Math. 3 (1998), 231-259.
- [26] M. Kontsevich, *Formal (non)commutative symplectic geometry*. The Gel'fand Mathematical Seminars, 1990-1992, 173-187, Birkhäuser Boston, Boston, MA, 1993.
- [27] M. Kontsevich and Y. Soibelman, *Notes on A_∞ algebras, A_∞ categories and non-commutative geometry*. I. 2004 (arXiv:math/0606241).
- [28] P. Lambrechts and D. Stanley, *Poincaré duality and commutative differential graded algebras*. Annales Scientifiques de l'École Normale Supérieure 41 (2008) 495-509.
- [29] J.-S. Loday, *Cyclic homology*. Second edition. Grundlehren der Mathematischen Wissenschaften, 301. Springer-Verlag, Berlin, 1998.
- [30] M. Penkava and A. Schwarz, *A_∞ algebras and the cohomology of moduli spaces*, in *Lie groups and Lie algebras: E. B. Dynkin's Seminar*, 91-107, Amer. Math. Soc. Transl. Ser. 2, 169, Amer. Math. Soc., Providence, RI, 1995.
- [31] D. Quillen, *Algebra cochains and cyclic cohomology*. IHES Publ. Math. No. 68 (1988), 139-174 (1989).
- [32] P. Seidel, *Vanishing cycles and mutation*, in *Proc. 3rd European Congress of Mathematics* (Barcelona, 2000), vol. II. Prog. Math., vol. 202, pp. 65-85. Basel: Birkhäuser 2001.
- [33] P. Seidel, *More about vanishing cycles and mutation*, in *Symplectic Geometry and Mirror Symmetry*. Proc. 4th KIAS International Conference (Seoul, 2000), pp. 429-465. Singapore: World Sci. 2001.
- [34] P. Seidel, *Fukaya categories and Picard-Lefschetz theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [35] T. Schedler, *A Hopf algebra quantizing a necklace Lie algebra canonically associated to a quiver*. Int. Math. Res. Not. 2005, no. 12, 725-760.
- [36] D. Sullivan, *String Topology: Background and Present State*, Current developments in mathematics, 2005, 41-88, Int. Press, Somerville, MA, 2007.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MI 48109,
UNITED STATES

E-mail address: xch@umich.edu